

FILE COPY

1

AD-A216 659

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE December 1989	3. REPORT TYPE AND DATES COVERED professional paper
4. TITLE AND SUBTITLE IMMUNODIAGNOSTIC APPROACHES TO DISEASES IN MARINE MAMMALS		5. FUNDING NUMBERS In-house	
6. AUTHOR(S) N. A. Vedros, J. P. Schroeder, and L. Suer		8. PERFORMING ORGANIZATION REPORT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Naval Ocean Systems Center San Diego, CA 92152-5000		10. SPONSORING/MONITORING AGENCY REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Naval Ocean Systems Center San Diego, CA 92152-5000		11. SUPPLEMENTARY NOTES	
12a. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution is unlimited.		12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) <p>The rapid diagnosis of infectious diseases in all animals has been the major focus of clinical microbiology for decades. It is particularly important in marine mammals since illness, with cessation of eating, causes rapid dehydration with serious sequelae.</p> <p>Initial studies indicated that commercial anti-Ig's of a variety of animals were not suitable for use in serological studies with marine mammals. The IgG and IgM of cetaceans and pinnipeds were isolated, antisera produced in rabbits and guinea pigs, and used to develop the single radial immunodiffusion (RID) immunoblot, and ELISA techniques. The base-line level of IgG and IgM in the bottlenose dolphin (<i>Tursiops truncatus</i>), the sea otter (<i>Enhydra lutris</i>), and the northern fur seal (<i>Callorhinus ursinus</i>) have been determined by RID. Specific antigens and monoclonal antibodies as positive controls for <i>Histoplasma</i>, <i>Blastomyces</i>, <i>Coccidioides</i>, <i>Candida</i>, and <i>Aspergillus</i> have been developed. Specific outer membrane proteins of <i>Erysipelothrix rhusiopathiae</i> and surface polysaccharides of <i>Pasturella multocida</i> and <i>Pseudomonas pseudomallei</i> have been used successfully in an ELISA assay. The potential of all of the above diagnostic tests for infectious diseases in marine mammals will be discussed.</p> <p>Published in <i>Proceedings First International Conference on Zoological and Avian Medicine</i>, September 1987.</p>			
14. SUBJECT TERMS mammals health		15. NUMBER OF PAGES	
17. SECURITY CLASSIFICATION OF REPORT UNCLASSIFIED		18. PRICE CODE	
18. SECURITY CLASSIFICATION OF THIS PAGE UNCLASSIFIED		19. SECURITY CLASSIFICATION OF ABSTRACT UNCLASSIFIED	
20. LIMITATION OF ABSTRACT UNLIMITED			

FROM: <input checked="" type="checkbox"/> TO: <input type="checkbox"/> BY: <input type="checkbox"/> DATE: <input type="checkbox"/>	
FROM: NOSC TO: <input type="checkbox"/> BY: <input type="checkbox"/> DATE: <input type="checkbox"/>	
FROM: <input type="checkbox"/> TO: <input type="checkbox"/> BY: <input type="checkbox"/> DATE: <input type="checkbox"/>	
FROM: <input type="checkbox"/> TO: <input type="checkbox"/> BY: <input type="checkbox"/> DATE: <input type="checkbox"/>	

PROCEEDINGS
FIRST INTERNATIONAL CONFERENCE
on
ZOOLOGICAL AND AVIAN MEDICINE



Sponsored by
Association of Avian Veterinarians
and
American Association of Zoo Veterinarians

September 6-11, 1987

Turtle Bay Hilton and Country Club
Oahu, Hawaii

Immunodiagnostic approaches to diseases in marine mammals. ¹Neylan A. vedros, Ph.D., ²J. P. Shroeder, and ¹L. Suer. ¹Biomedical and Environmental Health Sciences, University of California, Berkeley, CA 94720. ²NOSC Hawaii Laboratory, Kailua, Hawaii 96734.

The rapid diagnosis of infectious diseases in all animals has been the major focus of clinical microbiology for decades. It is particularly important in marine mammals since illness, with cessation of eating, causes rapid dehydration with serious sequelae.

Initial studies indicated that commercial anti-Ig's of a variety of animals were not suitable for use in serological studies with marine mammals. The IgG and IgM of cetaceans and pinnipeds were isolated, antisera produced in rabbits and guinea pigs, and used to develop the single radial immunodiffusion (RID) immunoblot, and ELISA techniques. The base-line level of IgG and IgM in the bottlenose dolphin (Tursiops truncatus), the sea otter (Enhydra lutris), and the northern fur seal (Callorhinus ursinus) have been determined by RID. Specific antigens and monoclonal antibodies as positive controls for Histoplasma, Blastomyces, Coccidioides, Candida, and Aspergillus have been developed. Specific outer membrane proteins of Erysipelothrix rhusiopathiae and surface polysaccharides of Pasturella multocida and Pseudomonas pseudomallei have been used successfully in an ELISA assay. The potential of all of the above diagnostic tests for infectious diseases in marine mammals will be discussed.